ORIGINAL ARTICLE

Test-retest Reliability of Unilateral and Bilateral Carrying **Protocol Based on Joule's Functional Capacity Evaluation**

Hanif Farhan Mohd Rasdi¹, Nurul Syahirah Md Senin²

¹ Centre Rehabilitation Sciences and Special Needs (iCaRehab), Occupational Therapy Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Jalan Raja Muda Abdul Aziz, Kuala Lumpur, Malaysia

² Flying Kids Occupational Therapy, No 6, Jalan horizon Perdana 1, Horizon Hills, 79150 Iskandar Puteri, Johor, Malaysia.

ABSTRACT

Introduction: Load carriage is one of the most frequent manual material handlings in industrial settings. However, due to the biomechanical risk exposures, carrying could lead to work-related musculoskeletal disorders. Joule's Functional Capacity Evaluation is an FCE system used widely in Malaysia in return-to-work programs to ensure work readiness and prevent re-injury. However, the literature regarding the test-retest reliability of the carrying protocols is minimal, leading to a questionable level of consistency. Therefore, this study aimed to investigate the test-retest reliability of the carrying protocols in Joule's FCE. Methods: A cross-sectional study was conducted to evaluate test-retest reliability where the carrying protocol of Joule's Functional Capacity Evaluation was used among healthy university students (N=30). The participants were asked to perform a retest of the carrying protocol after one week. Results: The ICC values for the carrying protocols were good for dominant unilateral carrying was (ICC: 0.82; 95% CI: -6.00 to 5.81), non-dominant carrying was (ICC: 0.74; 95% CI: -6.78 to 6.02) and bilateral carrying was (ICC: 0.85; 95% CI: -6.26 to 5.51). Bland and Altman's plot indicated no visible distribution patterns of the differences without exceeding the 95% limits of agreement. The standard error of measurements (SEMs) was relatively small for all carrying protocols. Conclusion: The test-retest reliability of carrying protocols in Joule's FCE was good. Future studies are needed to replicate this study in real patients to further verify the reliability for clinical settings.

Malaysian Journal of Medicine and Health Sciences (2022) 18(9):59-65.doi:10.47836/mjmhs18.s9.9

Keywords: Return-to-work, Load carriage, Work readiness, Ergonomics, Occupational therapy

Corresponding Author:

Hanif Farhan Mohd Rasdi, PhD Email: hanif_ot@ukm.edu.my Tel: 03-92897374

INTRODUCTION

Any accidents that cause injury, illness, or death in the workplace are considered occupational injuries (1). According to the literature, manual handling activities such as carrying and lifting are the main risk factors for waist soreness and various back disorders (2, 3). According to the Fourth European Working Conditions Survey (2005) carried out in the EU-27, 35% of all workers were exposed to carrying or moving heavy loads for at least a quarter of their working time (4). The factors that make manual handling hazardous are the load (e.g., too heavy, oversized, challenging to reach or grasp, unbalanced) and the task (e.g., too strenuous, involves awkward postures or movements) (5, 6).

The Social Security Organization (SOCSO) is a statutory body representing the employment accident insurance scheme and the invalidity pension scheme. This organization introduced a Return-to-Work (RTW) program in 2007, a comprehensive rehabilitation program for its insured persons with a disability due to work injury (7). When a worker has recovered from injury and is permitted to come back to work, this organization needs to assess the workers' level of work readiness. Therefore, a Functional Capacity Evaluation report is usually required to determine workers' physical capacity and disability (8).

Functional Capacity Evaluation (FCE) is a measurement tool that assists in determining safe, tolerable levels of function and predicting when an individual is ready to return to work duties (9). In addition, it also can provide a comprehensive assessment in determining the level of functionality of an individual in their work (10). Rehabilitation professionals frequently use performancebased outcome measures such as FCE to assess the level of functional disability since the measurements are more accurate and free from bias than self-report measures (9). There are various FCE types available, and one of them was Joule's FCE system by the Valpar International Corporation. Joule's FCE is a measurement tool that can evaluate an individual's functional abilities or limitations in safe, productive work tasks.

The FCE is considered the gold-standard assessment in a return-to-work program (11). The FCE has been practiced for years by occupational therapists, physiotherapists, vocational assessors and psychologists (9, 12). However, to the researcher's knowledge, the measurement properties, such as the validity and reliability of some of the FCE systems, are still limited. Generally, reliability can be defined as how an assessment tool can produce a stable and consistent decision (13). If a measurement can be replicated consistently within two or more time intervals, the tool is said to have good test-retest reliability (14). On the other hand, inaccurate FCE can increase the risk of re-injury. Therefore, professionals involved in work rehabilitation need to assess their patients' physical capabilities before returning to their previous work (15). According to the literature, the FCE had been criticized for lacking legitimacy, credibility and questionable outcomes unrelated to continuous work (16-18).

Thus, it is essential to determine and report the validity and reliability of the FCE to allow accurate clinical decision-making. For instance, in the RTW program established by the SOCSO, the FCE report is critical to determine the ability of a client to return to work and for legal documentation purposes. A study has shown that intra-rater reliability between the assessors for Joule's FCE is high in healthy adults (10), but no study investigating the reliability in Malaysia. Therefore, the purpose of this study was to examine the test-retest reliability of the carrying protocol in Joule's FCE.

MATERIALS AND METHODS

Study design

This cross-sectional study was conducted on healthy university students recruited using convenience sampling (n=33). The initial sample consisted of 33 participants: 16 males and 17 females. However, only 30 participants performed the retest (13 males and 17 females). All declared to be healthy, i.e., having no medical condition that would restrict them from performing maximally. The data were collected within three months.

Sample size

The sample size determination for this study was conducted based on the power analysis method. The power set for this study was 80%, while the alpha value was set at 5%. The effect size used for this sample size

calculation was based on the value of the Intra-class Correlation Coefficient (ICC) of 0.87 (19, 20). After considering 10% of dropout rates, 30 participants were required in this study.

Instruments

This study used two instruments, namely the Joule's Functional Capacity Evaluation (FCE) (Valpar International Corporation) and the International Physical Activity Questionnaire Short Form (IPAQ-SF). Joule's FCE was selected due to the limited literature about this FCE. This instrument can be used to assess the ability of a person to function in manual handling activities and demonstrated high intra-rater reliability (ICC > 0.90) for the Joule's FCE in healthy adults (10). The IPAQ-SF was used to determine the level of physical activity in the last seven days (i.e., inactive, moderately active or active). This questionnaire was selected due to the ease of assessing participants' physical activity over the last 7-days. The IPAQ-SF consisted of seven items, including open-ended questions about the physical activity of the previous seven days. It is a self-assessment report, paperpencil version or verbally. The correlation coefficients of the IPAQ-SF show ranged from 0.16 to 0.35 indicating moderate criterion validity, while (ICC = 0.71-0.89) indicated acceptable reliability (21, 22).

Procedures

The main parameters of this study are the carrying load limit for unilateral and bilateral carrying protocols in Joule's FCE. The carrying protocols were conducted based on the standardized methods as stated in the Joule FCE module. The participants were introduced to the general FCE procedures for carrying protocol and then signed informed consent. Then, they were asked to fill in the IPAQ-SF and performed the carrying protocol with progressive loads (i.e., 1.5 kg increment). The loads were placed into standard containers with dividers. The load arrangement in the dividers was based on the Joule's load progression flipchart. The participants were briefly instructed on the required performance. The evaluator demonstrated each test of carrying protocol (i.e., unilateral and bilateral carrying). After a one week, carrying protocols were retested. For the unilateral carrying activity, the participants lifted a container using the dominant hand, carried it from point A to point B (7 meters), and returned to point A. After that, the participants changed to their non-dominant hands, took the load from point A to point B, and returned to point A. For the bilateral carrying activity, the participants lifted the container using both hands, carrying the load from point A to point B and returning to point A. Finally, each test was terminated for three reasons: 1) the participants wished to do so, 2) heart rate exceeded 85% age-related target heart rate, and 3) unsafe action, defined as a situation in which the participants were not in complete control of him or herself or the load. The evaluator recorded the time and safe maximum carrying load limit directly after each test.

Data Analysis

The data were analyzed using IBM SPSS Statistics version 22. The sociodemographic characteristics and level of physical activity were analyzed using The Intra-class Correlation descriptive analysis. Coefficient (ICC) was used to determine the level of agreement between the measurements. An ICC greater than 0.7 was considered good, and an ICC over 0.9 was excellent (23). The Bland-Altman (B&A) plot was used to determine the 95% limit of agreement between the measurements. Bland and Altman introduced a method to describe the limits of agreement between two quantitative measurements (24). They had established a way to quantify the agreement between two quantitative measures by constructing (25) limits of agreement. These statistical limits are calculated using the mean and the standard deviation (s) of the differences between the two measurements. Bland and Altman recommended that 95% of the data points lie within \pm 2s of the mean difference (26). Finally, the following formula was used to determine the standard error of measurement (SEM), where the SD (diff) is the standard deviation of the difference between the measurements, and the ICC is the value of the Intraclass Correlation

Coefficient: *SD* (*diff*) $\sqrt{1 - ICC}$

Ethical Clearance

The ethical clearance was obtained from the UKM Secretariat of Ethics (ethics numbers: NN-2017-187).

RESULT

Thirty-three participants met the inclusion criteria, and all returned the consent forms. However, only 30 participants return on retest (13 males & 17 females). The descriptions of the 30 participants are presented in Table I. The means, standard deviations, ICCs and standard error of measurement (SEM) are shown in Table 2. The ICC values of the dominant hand, non-dominant hand and bilateral carrying task were 0.82, 0.74 and 0.85, respectively. The standard error of measurement (SEM) was relatively minor for all types of carrying (SEM_{dominant} was 1.27kg; SEM_{non-dominant} = 1.678; SEM_{bilateral} = 1.15) (Table II).

Bland and Altman's plot indicated that the dominant hand's carrying had an outlier beyond the upper limit and the lower limit for the limits of agreement. For the nondominant hand, Bland and Altman's plot indicated two outliers above the upper limit, while bilateral carrying had one outlier beyond the lower limit. The differences show a heteroscedasticity pattern for dominant unilateral. However, non-dominant unilateral and bilateral carrying shows a linear pattern of differences. In addition, the mean differences for dominant and non-dominant unilateral and bilateral carrying were -0.0933, -0.3833 and -0.3767, respectively, which are relatively close to zero, indicating a good level of agreement.

Fable I: Participants	' characteristics	(N=30)
-----------------------	-------------------	--------

I		•	
	Freq. (%)	Mean (SD)	
Gender			
Female	17 (57%)		
Male	13 (43%)		
Age			
20	2 (7%)		
21	6 (20%)	22.60 (1.248)	
23	17 (57%)		
24	4 (13%)		
25	1 (3%)		
Race			
Malay	16 (53%)		
Chinese	12 (40%)		
Indian	2 (7%)		
First IPAQ			
Inactive	8 (27%)	1.02 (0.6011)	
Minimal active	16 (53%)	1.93 (0.0911)	
HEPA active	6 (20%)		
Second IPAQ			
Inactive	3 (10%)	2.10 (0.548)	
Minimal active	21 (70%)		
HEPA active	6 (20%)		

Table II: Test-retest reliability of carrying task Joule's Functional Capacity Evaluation

Type of carrying	ICC ·	95% Cl		CENA
		Lower	Upper	SEM
Dominant-hand unilateral carrying	0.82	-6.00	5.81	1.27
Non-dominant unilateral carrying	0.74	-6.78	6.02	1.68
Bilateral carrying	0.85	-6.26	5.51	1.15

*ICC = Intra-Class Correlation Coefficient. SEM = Standard Error of Measurement

DISCUSSION

The findings from this study indicated that the testretest reliability of the carrying protocol in Joule's FCE was acceptable, indicating that the carrying protocols are consistent from one testing occasion to another. In addition, the findings were consistent with a previous study that reported good test-retest reliability of carrying task (25). For unilateral carrying, the dominant hand indicated better reliability than non-dominant unilateral carrying, indicating a slight difference in reliability between both unilateral carrying protocols. Reneman, Brouwer (27) also reported differences in the reliability score between unilateral carrying. Furthermore, when comparing between types of carrying, bilateral load carriage showed a higher level of agreement than both unilateral carrying protocols. This finding may suggest that the bilateral carrying was relatively stable compared with unilateral carrying. Carrying loaded bilaterally will displace one's centre of gravity posteriorly. Hence, increasing an individual's pelvic tilt or forward lean will keep them in an upright, vertical position (28-30). Even if the total load weight doubles during bilateral carrying, it does not increase the spine load (31). However, when observing the biomechanics of unilateral carrying, the trunk bends towards the contralateral side (unload side) to compensate for the perturbation from the external load (32). This action causes higher forces toward the spinal while walking, leading to increased muscular activity and more significant spinal shear and compressive force (30, 33). Furthermore, continuously carrying a load on one side of the body can lead to muscle fatigue due to increased muscle use on opposite sides since carrying one hand results in high spine compression when bilateral carrying (34). This fatigue phenomenon can affect the consistency of the result and then influence the reliability of this unilateral carrying task (35). Nevertheless, bilateral carrying becomes the most demanding task compared with other carrying on the worksite (36). Hence, it is vital to ensure this carrying is stable enough as an ergonomic intervention to prevent injury on the worksite.

According to the limits of agreement, most of the differences were within the upper and lower limit of agreement. However, there were some outliers in both types of carrying protocols. Two participants (outliers) in the dominant and non-dominant unilateral carrying, and one participant scored outside of the 95% limits of agreement. Based on the level of physical activity before the test and retest, all the participants had no changes in their physical activity level. However, slight differences in Metabolic Equivalents (METs) were indicated, either increase or decrease. Dodds, Kuh (37) stated that increasing physical activity benefits grip strength, while Stenholm, Tiainen (38) stated becoming sedentary was associated with a decline in muscle strength. Hence, this evidence correlated with participants carrying a heavier load on retest when increasing METs before the retest. However, this situation is not associated with one participant (ID=29) who had a heavier load while decreasing METs before the retest.

Pre-existing fatigue during the retest session can also explain the differences between carrying activities. It isn't easy to maintain physical activity as students since they have to participate in numerous co-curricular activities. Therefore, prolonged physical activity between the tests may lead to fatigue. A previous study stated intensive training of muscles causes a decline in performance (35). For instance, Ahmed (39) suggested avoiding upper extremity fatigue among basketball players since it may reduce grip strength and affect performance. Fatigue has two types which are central and peripheral fatigue. During the prolonged exercise of moderate intensity, a decrease in the blood glucose level due to depletion of the liver glycogen stores is one factor known to affect the central nervous system and cause fatigue (40). Another factor that can cause central fatigue during dynamic exercise is the increased neurotransmitter release in the brain (41-43). However, such adaptations may contribute to the delay of central fatigue during sustained exercise. This fatigue indicates a reduction in the contractile strength of the muscle fibres and changes in the mechanisms underlying the transmission of muscle action potentials (44).

This study's standard error of measurements (SEMs) was relatively small for all carrying protocols, suggesting a clinically tolerable measurement error for the evaluation. Bilateral carrying indicated the smallest SEM rather than unilateral carrying, indicating better quality, accuracy, and reliability (45). Even though the SEM of unilateral carrying was slightly higher than bilateral carrying, it can still be acceptable since all measurements contain some error. No similar SEM data for FCE carrying tests have been previously reported. Thus, the SEM's cut-off value for clinical tolerance cannot be retrieved from literature and should be based on the practitioner's knowledge of the carrying tests. However, compared with the SEM for other functional assessments, the study also shows relatively small SEM within 0.19 to 0.66 with high reliability of measurements (46). In addition, the SEM lifting tests varied between 1.9 and 8.6 kg, and the SEM for carrying lifting strength test was 3.4 (47). Thus, this study indicated a sufficient level of agreement with the lifting tests.

To our knowledge, this is the first study that has investigated the test-retest reliability of the carrying protocol of Joule's FCE in Malaysia. This study has found that the carrying protocol of Joule's FCE has good testretest reliability. Therefore, it is essential to ensure this instrument is safe, reliable, valid, practical, and utility (12, 43). However, some difficulties were faced in carrying out this study, such as setting up participants' appointments for their retest sessions. In addition, most of them were university students with busy schedules due to classes and co-curricular activities. In addition, the time to access the instrument was limited as it was located in a teaching clinic. Therefore, it can only be accessed when no patients or classes are being held in the room. However, despite these challenges, an adequate sample size was successfully achieved.

Although the number of participants in this study achieved a sample size considered adequate for a fair quality study, a larger sample size would have improved the quality of the study results (48). Besides, this study was conducted among healthy university students, which can pose different effects on retest due to several factors such as differences in lifestyles. However, the influence of physical activity on the carrying load limit was not investigated in detail throughout the study due to the statistical limitations. Therefore, future studies are needed to improve the statistical power and replicate this study in clinical situations to verify the reliability further.

CONCLUSION

The primary purpose of this research was to investigate the test-retest reliability for the carrying protocol of Joule's FCE among healthy adults. The findings show the carrying protocol in Joule's FCE was good testretest reliability. Since the carrying protocol in Joule's FCE is reliable, it can help clinicians make decisions in clinical settings to prevent re-injury among the workers. As this study was conducted on healthy participants, future studies are needed to replicate this study in clinical situations such as work-related musculoskeletal disorders (WRMD) to verify the reliability.

ACKNOWLEDGEMENTS

Much appreciation is given to Mr. Lim Wen Xing for helping in the analyzed data collection and member checking process. Many thanks to all participants, students at the Universiti Kebangsaan Malaysia (Kuala Lumpur Campus), for participating in this study.

REFERENCES

- 1. Spector J.T., Masuda Y.J., Wolff N.H., Calkins M., & Seixas N. Heat exposure and occupational injuries: review of the literature and implications. Current environmental health reports. 2019;6(4):286-96. DOI: 10.1007/s40572-019-00250-8
- 2. Kędra A., Plandowska M., Kędra P., & Czaprowski D. Non-specific low back pain: cross-sectional study of 11,423 children and youth and the association with the perception of heaviness in carrying of schoolbags. PeerJ. 2021;9:e11220. https://doi.org/10.7717/peerj.11220
- 3. Suehiro T., Ishida H., Kobara K., Osaka H., & Watanabe S. Altered trunk muscle recruitment patterns during lifting in individuals in remission from recurrent low back pain. Journal of Electromyography and Kinesiology. 2018;39:128-33. DOI: 10.1016/j.jelekin.2018.02.008
- 4. Parent-Thirion A., Fernández Macías E., Hurley J., & Vermeylen G. Fourth European working conditions survey, European foundation for the improvement of living and working conditions. Luxembourg: Office for Official Publications of the European Communities. 2007.
- 5. Das B. Effects of Awkward Posture on Work-Related Musculoskeletal Disorders (WMSDs) among Sawmill Workers in India. Journal of Occupational Health and Epidemiology. 2020;9(3):158-66.

DOI:10.29252/johe.9.3.158

- 6. Ramandi F.F., Study of low back pain intensity and disability index among manual material handling workers of a tile and ceramic industrial unit, Iran (2016). Journal of Occupational Health and Epidemiology. 2018;7(3):167-73. 10.29252/ johe.7.3.167
- 7. Mohammed M,. The Return to Work Programme in Malaysia-investing in people. International Journal of Disability Management. 2014;9. DOI:10.1017/ IDM.2014.8
- Sinden, K. E., McGillivary, T. L., Chapman, E., & Fischer, S. L. Survey of kinesiologists' functional capacity evaluation practice in Canada. Work (Reading, Mass.), 2017;56(4), 571–580. https://doi. org/10.3233/WOR-172519
- De Baets, S., Calders, P., Schalley, N., Vermeulen, K., Vertriest, S., Van Peteghem, L., Coussens, M., Malfait, F., Vanderstraeten, G., Van Hove, G., & Van de Velde, D. Updating the Evidence on Functional Capacity Evaluation Methods: A Systematic Review. Journal of occupational rehabilitation,2018;28(3),418–428.https://doi. org/10.1007/s10926-017-9734-x
- 10. Mitchell, D., Hancock, E., & Alexander, L. An investigation of the inter-rater reliability of the Valpar Joule functional capacity evaluation in healthy adults. Work (Reading, Mass.), 2015;53(2), 337–345. https://doi.org/10.3233/WOR-152154
- 11. McFadden, S., MacDonald, A., Fogarty, A., Le, S., & Merritt, B. K. Vocational assessment: a review of the literature from an occupationbased perspective. Scandinavian journal of occupational therapy. 2010 ;17(1), 43–48. https:// doi.org/10.1080/11038120903096633
- Hart, D. L., Isernhagen, S. J., & Matheson, L. N.
 Guidelines for functional capacity evaluation of people with medical conditions. The Journal of orthopaedic and sports physical therapy. 1993;18(6), 682–686. https://doi.org/10.2519/ jospt.1993.18.6.682
- 13. Phelan C, Wren J. Reliability is the degree to which an assessment tool produces stable and consistent results. Assessment. 2005:06.
- 14. Scheel C, Mecham J., Zuccarello V., & Mattes R. An evaluation of the inter-rater and intrarater reliability of OccuPro's functional capacity evaluation. Work. 2018;60(3):465-73. https://doi. org/10.3233/WOR-182754
- 15. Genovese E, Galper JS. Guide to the evaluation of functional ability: how to request, interpret, and apply functional capacity evaluations: American Medical Association; 2009. DOI:10.1007/s10926-010-9279-8
- 16. Gouttebarge V., Kuijer P.P.F., Wind H., van Duivenbooden C., Sluiter J.K., & Frings-Dresen M.H. Criterion-related validity of functional capacity evaluation lifting tests on future work disability risk and return to work in the construction

industry. Occupational and environmental medicine. 2009;66(10):657-63. doi:10.1136/ oem.2008.042903

- 17. Gross, D. P., & Batti*é*, M. C. Does functional capacity evaluation predict recovery in workers' compensation claimants with upper extremity disorders? Occupational and environmental medicine. 2006;63(6):404-10. https://doi.org/10.1136/oem.2005.020446
- 18. Gross D.P., Battié M.C., Cassidy J.D. The prognostic value of functional capacity evaluation in patients with chronic low back pain: part 1: timely return to work. Spine. 2004;29(8):914-9. DOI: 10.1097/00007632-200404150-00019
- 19. Walter S., Eliasziw M., & Donner Α., Sample optimal designs size and for Statistics in reliability studies. medicine. 1998;17(1):101-10. https://doi.org/10.1002/ (sici)1097-0258(19980115)17:1<101::aidsim727>3.0.co;2-e
- 20. Winer B. Statistical Principles in Experimental Design: 2d Ed: McGraw-Hill; 1971.
- 21. Dinger M.K, Behrens T.K., & Han J.L. Validity and reliability of the International Physical Activity Questionnaire in college students. American Journal of Health Education. 2006;37(6):337-43. DOI: 10.1080/19325037.2006.10598924
- Ekelund, U., Sepp, H., Brage, S., Becker, W., Jakes, R., Hennings, M., & Wareham, N. J. Criterionrelated validity of the last 7-day, short form of the International Physical Activity Questionnaire in Swedish adults. Public health nutrition. 2006;9(2):258-65.https://doi.org/10.1079/ phn2005840
- 23. Liljequist D, Elfving B, Skavberg Roaldsen K. Intraclass correlation–A discussion and demonstration of basic features. PloS one. 2019;14(7):e0219854.
- 24. Bland J.M., & Altman D.G. Measuring agreement in method comparison studies. Statistical methods in medical research. 1999;8(2):135-60. doi: 10.1177/096228029900800204
- 25. Branton E.N., Arnold K.M., Appelt S.R., Hodges M.M., Battié M.C., & Gross D.P. A short-form functional capacity evaluation predicts time to recovery but not sustained return-to-work. Journal of occupational rehabilitation. 2010;20(3):387-93. DOI: 10.1007/s10926-010-9233-9
- 26. Giavarina D. Understanding bland altman analysis. Biochemia medica: Biochemia medica. 2015;25(2):141-151. DOI: https://doi. org/10.11613/BM.2015.015
- Reneman, M. F., Brouwer, S., Meinema, A., Dijkstra, P. U., Geertzen, J. H., & Groothoff, J. W. . Test-retest reliability of the Isernhagen work systems functional capacity evaluation in healthy adults. Journal of Occupational Rehabilitation. 2004;14(4):295-305. https://doi.org/10.1023/ b:joor.0000047431.40598.47

- 28. Pascoe, D. D., Pascoe, D. E., Wang, Y. T., Shim, D. M., & Kim, C. K. Influence of carrying book bags on gait cycle and posture of youths. Ergonomics. 1997;40(6):631-640. DOI: 10.1080/001401397187928
- 29. Smith, B., Ashton, K. M., Bohl, D., Clark, R. C., Metheny, J. B., & Klassen, S. Influence of carrying a backpack on pelvic tilt, rotation, and obliquity in female college students. Gait & posture. 2006;23(3):263-267. https://doi.org/10.1016/j. gaitpost.2005.02.009
- Hyung, E., Lee†, H., & Kwon, Y. Influence of load and carrying method on gait, specifically pelvic movement. Journal of physical therapy science. 2016;28(7):2059-2062. DOI:10.1589/jpts.28.2059
- 31. McGill, S. M., Marshall, L., & Andersen, J. Low back loads while walking and carrying: comparing the load carried in one hand or in both hands. Ergonomics. 2013;56(2):293-302. https://doi.org/1 0.1080/00140139.2012.752528
- 32. Zhang, X. A., Ye, M., & Wang, C. T. Effect of unilateral load carriage on postures and gait symmetry in ground reaction force during walking. Computer methods in biomechanics and biomedical engineering. 2010;13(3):339-44. https://doi.org/10.1080/10255840903213445
- Bordelon, N. M., Wasserberger, K. W., Cassidy, M. M., & Oliver, G. D. The Effects of Load Magnitude and Carry Position on Lumbopelvic-Hip Complex and Scapular Stabilizer Muscle Activation During Unilateral Dumbbell Carries. The Journal of Strength & Conditioning Research. 2021;35:S114-S9. DOI: 10.1519/JSC.00000000003880
- 34. Corrigan L. The effect of unilateral load carriage on the muscle activities of the trunk and lower limbs of young healthy males during gait: University of Ottawa (Canada); 2012. http://dx.doi. org/10.20381/ruor-6220
- 35. Pasiakos, S. M., Lieberman, H. R., & McLellan, T. M. Effects of protein supplements on muscle damage, soreness and recovery of muscle function and physical performance: a systematic review. Sports Medicine. 2014;44(5):655-70. https://doi. org/10.1007/s40279-013-0137-7
- 36. Amin Z, Mohammad R, Azizc SA, Othmand N. Workers' Safety Awareness Level on Hand Related Injury Accident in Metal Fabrication Industry. Journal of Advanced Research in Applied Sciences and Engineering Technology. 2015;1:1-12.
- 37. Dodds, R., Kuh, D., Aihie Sayer, A., & Cooper, R. Physical activity levels across adult life and grip strength in early old age: updating findings from a British birth cohort. Age and ageing. 2013;42(6):794-8.https://doi.org/10.1093/ageing/ aft124
- Stenholm, S., Tiainen, K., Rantanen, T., Sainio, P., Heliuvaara, M., Impivaara, O., & Koskinen, S. Long - termdeterminantsofmusclestrengthdecline: Prospective evidence from the 22 - year mini -

Finland follow - up survey. Journal of the American Geriatrics Society. 2012;60(1):77-85. https:// doi.org/10.1111/j.1532-5415.2011.03779.x DOI:10.2478/hukin-2013-0027

- 39. Ahmed T. The effect of upper extremity fatigue on grip strength and passing accuracy in junior basketball players. Journal of human kinetics. 2013;37(1):71-9.
- Nybo, L., Møller, K., Pedersen, B. K., Nielsen, B., & Secher, N. H. Association between fatigue and failure to preserve cerebral energy turnover during prolonged exercise. Acta Physiologica. 2003;179(1):67-74. https://doi.org/10.1046/ j.1365-201X.2003.01175.x
- 41. Meeusen, R., & Roelands, B. Central fatigue and neurotransmitters, can thermoregulation be manipulated? Scandinavian journal of medicine & science in sports. 2010;20:19-28. https://doi. org/10.1111/j.1600-0838.2010.01205.x
- 42. Meeusen, R., & Roelands, B. Fatigue: is it all neurochemistry? European journal of sport science. 2018;18(1):37-46. https://doi.org/10.1080/174613 91.2017.1296890
- 43. Trippolini, M. A., Dijkstra, P. U., Jansen, B., Oesch, P., Geertzen, J. H., & Reneman, M. F. Reliability of clinician rated physical effort determination during functional capacity evaluation in patients with chronic musculoskeletal pain. Journal of occupational rehabilitation. 2014;24(2):361-9. doi: 10.1007/s10926-013-9470-9
- 44. Boyas, S., & Guével, A. Neuromuscular fatigue in healthy muscle: underlying factors and adaptation

mechanisms. Annals of physical and rehabilitation medicine. 2011;54(2):88-108. https://doi. org/10.1016/j.rehab.2011.01.001

- 45. Tighe, J., McManus, I., Dewhurst, N.G. et al. The standard error of measurement is a more appropriate measure of quality for postgraduate medical assessments than is reliability: an analysis of MRCP (UK) examinations. BMC medical education. 2010;10(1):40. https://doi.org/10.1186/1472-6920-10-40
- 46. Srisim, K., Saengsuwan, J., & Amatachaya, S. Functional assessments for predicting a risk of multiple falls in independent ambulatory patients with spinal cord injury. The journal of spinal cord medicine. 2015;38(4):439-45. https://doi.org/10.1 179/2045772313Y.0000000186
- 47. Gouttebarge V., Wind H., Kuijer P.P., Sluiter J.K., & Frings-Dresen M.H. Reliability and agreement of 5 Ergo-Kit functional capacity evaluation lifting tests in subjects with low back pain. Archives of physical medicine and rehabilitation. 2006;87(10):1365-70. https://doi.org/10.1016/j.apmr.2006.05.028
- Terwee, C. B., Mokkink, L. B., Knol, D. L., Ostelo, R. W., Bouter, L. M., & de Vet, H. C. Rating the methodological quality in systematic reviews of studies on measurement properties: a scoring system for the COSMIN checklist. Quality of Life Research. 2012;21(4):651-7. https://doi. org/10.1007/s11136-011-9960-1